

Experimental test of Hatano and Sasa's nonequilibrium steady-state equality

E. H. Trepagnier^{*†}, C. Jarzynski[‡], F. Ritort[§], G. E. Crooks[¶], C. J. Bustamante^{*†||**}, and J. Liphardt^{*†||††}

^{*}Biophysics Graduate Group and Departments of [¶]Physics, ^{**}Molecular and Cell Biology, and [¶]Plant and Microbial Biology, University of California, Berkeley, CA 94720; [‡]Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545; [§]Departament de Física Fonamental, Universitat de Barcelona, Barcelona 08028, Spain; and [†]Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

Contributed by C. J. Bustamante, September 3, 2004

Most natural processes occur far from equilibrium and cannot be treated within the framework of classical thermodynamics. In 1998, Oono and Paniconi [Oono, Y. & Paniconi, M. (1998) *Prog. Theor. Phys. Suppl.* 130, 29–44] proposed a general phenomenological framework, steady-state thermodynamics, encompassing nonequilibrium steady states and transitions between such states. In 2001, Hatano and Sasa [Hatano, T. & Sasa, S. (2001) *Phys. Rev. Lett.* 86, 3463–3466] derived a testable prediction of this theory. Specifically, they were able to show that the exponential average of Y , a quantity similar to a dissipated work, should be equal to zero for arbitrary transitions between nonequilibrium steady states, $-\ln\langle e^{-Y} \rangle = 0$. We have tested this strong prediction by measuring the dissipation and fluctuations of microspheres optically driven through water. We have found that $-\ln\langle e^{-Y} \rangle \approx 0$ for three different nonequilibrium systems, supporting Hatano and Sasa's proposed extension of thermodynamics to arbitrary steady states and irreversible transitions.

Classical thermodynamics and statistical mechanics are organized around the concept of “equilibrium states.” However, most processes of interest, such as fast switching between phonon distributions in optical cavities (1), oscillations of pumped mesoscopic chemical reaction systems (2, 3), and biological reactions by molecular machines (4), occur far from equilibrium. Molecular pumps, for example, maintain ions at nonequilibrium concentrations by active transport across membranes, and intermediate metabolism in cells involves transitions between nonequilibrium steady states for which there is no thermodynamic description.

Part of the difficulty in formulating a general theory of nonequilibrium thermodynamics is the diversity of dissipation regimes and mechanisms in systems away from thermal equilibrium. With this in mind, the study of “nonequilibrium steady states” occupies an interesting middle ground between the familiar territory of equilibrium thermodynamics and a nonequilibrium “thermodynamics of everything.” In fact, several theories have been proposed in which the state space includes such nonequilibrium steady states, which are characterized by the flow of constant currents (of energy, mass, or charge), as well as equilibrium states, which are defined by the absence of currents (5–8).

Any extension of thermodynamics should include a generalization of the second law. In one of its several classic formulations, this law places a constraint on transitions between equilibrium states, expressed by the Clausius inequality (9),

$$\int_A^B \frac{dQ}{T} \leq \Delta S, \quad [1]$$

where the left side pertains to a transition from state A to state B , and the right side is the entropy difference between these states. Does a similarly universal law govern transitions between nonequilibrium steady states?

One can address this question at the level of macroscopic phenomenology (see ref. 8), or one can seek the answer in a

microscopic, statistical analysis. Taking the latter approach, Keizer (5, 10) derived a generalized Clausius inequality by considering the steady-state fluctuations of a set of extensive variables $\mathbf{n} = (n_1, n_2, \dots)$, which provide a coarse-grained snapshot of the system. This result describes linear-response behavior in the near-steady-state regime.

More recently, Hatano and Sasa (11), motivated by the phenomenological framework of Oono and Paniconi (8), have obtained predictions for transitions between steady states of a system expressed in terms of the evolution of its microscopic degrees of freedom. Their results remain valid even if the system is driven away from steady-state behavior, suggesting that a strong Clausius-like inequality for transitions between nonequilibrium steady states may exist. Specifically, they predict that a Boltzmann-weighted average of the transition dissipation is equal to zero, such that $-\ln\langle e^{-Y} \rangle = 0$. Here, we report experiments in which optically dragged microspheres are used to test the predictions of Hatano and Sasa (11), which we now introduce.

Theory and Background

Let x denote the microstate of some system of interest, let α denote an externally controlled parameter, and let us suppose that, when this parameter is held fixed, the system relaxes to a stationary state described by a probability distribution $\rho_{ss}(x; \alpha)$. Here, we use the term “stationary state” quite generally, referring to either an equilibrium state or a nonequilibrium steady state. In the former case, ρ_{ss} is the familiar Boltzmann–Gibbs distribution; in the latter case, ρ_{ss} describes the microscopic fluctuations of the system in the steady state. When we carry the system from one stationary state to another, its microscopic evolution is specified by a trajectory $x(t)$.

Imagine an idealized process by which we drive the system from one stationary state to another by holding the parameter fixed at α_1 and then varying it over a finite time τ to a new constant value α_2 . If we change the parameter slowly and gently, the system moves through a continuous sequence of stationary states and the dissipated work is at a minimum. If we instead vary the parameter rapidly and violently, then the system cannot relax to the state consistent with the current parameter value because that value is changing too quickly. For transitions between equilibrium states, the Clausius inequality provides a quantitative measure of this lag: the more irreversible the process, the greater the difference between the two sides of Eq. 1. For transitions between nonequilibrium steady states, Hatano and Sasa (11) identified a new property, the Y value, which measures this lag. They construct the following quantity:

$$Y = \int_0^\tau dt \, \dot{\alpha}(t) \frac{\partial \phi}{\partial \alpha}(x(t), \alpha(t)), \quad [2]$$

^{††}To whom correspondence should be addressed. E-mail: liphardt@physics.berkeley.edu.

© 2004 by The National Academy of Sciences of the USA

where $\phi(x, \alpha) = -\ln \rho_{ss}(x; \alpha)$ and $\dot{\alpha} = d\alpha/dt$. The value of Y depends on both what we do to the system, as specified by the imposed time-dependence of the parameter $\alpha(t)$, and on how the system responds, which is represented by the phase-space trajectory $x(t)$. Because our system is subject to thermal noise, each repetition of the process yields a different Y value. Now, consider a statistical ensemble of trajectories, obtained by repeatedly varying the control parameter according to the same schedule $\alpha(t)$. Under very general conditions, Hatano and Sasa have shown that:

$$\langle e^{-Y} \rangle = 1, \quad [3]$$

where the angular brackets denote an average over our ensemble of repetitions of the process. By Jensen's inequality (12), Eq. 3 implies that:

$$\langle Y \rangle \geq 0. \quad [4]$$

Although the analysis of Hatano and Sasa (11) was carried out in the specific context of a trapped Brownian particle subject to a nonconservative force, it is evident from their derivation of Eqs. 3 and 4 that they are more general and do not depend on particular assumptions about the dynamics of the system.

Connecting their results to earlier work by Oono and Paniconi (8), Hatano and Sasa (11) interpret Eq. 4 as a generalized second law of thermodynamics that is applicable to transitions between (equilibrium or nonequilibrium) stationary states. As explained in greater detail in ref. 11, several observations support this interpretation. First, $Y = 0$ when the process is carried out reversibly, suggesting that in the more general case the non-negative value $\langle Y \rangle$ provides a measure of the irreversibility of the process. Second, for the system studied in ref. 11, Eq. 4 is equivalent to a generalized Clausius inequality proposed within Oono and Paniconi's phenomenological steady-state thermodynamic framework (8). Finally, Eqs. 3 and 4 reduce to known results for transitions between equilibrium states in the appropriate limit (see *Results and Discussion*).

To test Hatano and Sasa's predictions, we dragged a microscopic bead through water by using a steerable harmonic optical trap. Our protocol created nonequilibrium steady states that are both experimentally tractable and theoretically understood. On the experimental side, laser tweezers and similar optical devices have proven to be ideal laboratory tools for probing nonequilibrium statistical physics at microscopic length scales (13–17). In the present work, this technology provided the means to pull the bead and also to observe its motion at the level of thermal fluctuations. On the theoretical side, treating the bead as a Brownian particle and the optical trap as a harmonic potential, we easily obtain an expression for the steady-state distribution, ρ_{ss} , which enters into the definition of Y in Eq. 2. We subsequently confirmed the validity of the theoretical predictions for ρ_{ss} of Mazonka and Jarzynski (18) by comparing them with the experimentally measured distributions, and we found excellent agreement (data not shown). We chose a system in which ρ_{ss} is known *a priori*. In fact, Hatano and Sasa's relation for transitions between steady states can be applied to any physical system in which the stationary distribution ρ_{ss} is known or can be extracted from experiments. The first step in applying Hatano and Sasa's relation to more complicated systems (for example, in turbulence or granular media) in which the steady-state distributions ρ_{ss} are not known, will be to obtain them experimentally.

Methods

For each experiment, micron-sized polystyrene beads (diameter, 10.06 μm ; Bangs Laboratories, Carmel, IN) were dispensed and diluted inside a microfluidics chamber, which was then sealed. A single bead was then trapped in a harmonic potential created by




Fig. 1. Schematic representation of the experiment. Optically trapped bead lags a distance x behind the center of the trap translating at velocity v .

superposing the foci of two counterpropagating 834-nm laser beams (19), and an ultra-fast steerable mirror (Nano MTA-2, Mad City Labs, Madison, WI) was used to translate the trap. The optical force $f(t)$ exerted by the trap is equal in magnitude and opposite in direction to the rate of change of the momentum of light (19), which we measured directly by using position-sensitive photodetectors (DL-10, United Detector Technology Sensors). The position of the trap was calculated from the angular rotation of the steerable mirror. Data were sampled at 10 kHz, with the average over 10 data points saved to a disk. The following further parameters characterized the experiments: the trap constant $\kappa[pN/\mu\text{m}]$; the number of switching repetitions N ; the trap velocities v_1 and v_2 ($\mu\text{m/s}$), corresponding to the initial and final steady states, respectively; the switching time τ (s); and the value of $q = \kappa/\beta\gamma[pN(\mu\text{m/s})]$. Values for the first experiment were as follows: $\kappa = 4.25$, $n = 3,924$, $v_1 = 8.12$, $v_2 = 12.15$, $\tau = 0.06$, and $q = 0.20$. Values for the second experiment were as follows: $\kappa = 4.51$, $n = 3,163$, $v_1 = 9.93$, $v_2 = 13.56$, $\tau = 0.06$, and $q = 0.21$. Values for the third experiment were as follows: $\kappa = 4.9$, $n = 3,603$, $v_1 = 7.53$, $v_2 = 10.20$, $\tau = 0.08$, and $q = 0.23$.

Results and Discussion

Nonequilibrium steady states were created by translating the optical trap at constant speed (Fig. 1). After a relaxation time $t_R \sim \gamma/\kappa$, where γ is the friction coefficient of the bead in solution and κ is the spring constant of the trap, the bead settled into a steady state in which its position fluctuated around an average displacement $\gamma v/\kappa$ behind the minimum of the trap potential, so that the average force exerted by the trap balanced the average frictional force felt by the bead. This nonequilibrium steady state was maintained by a continual transfer of energy: the trap performed work on the bead at an average rate $\dot{P} = \gamma v^2$, which was dissipated as heat into the surrounding buffer.

We began our experimental trajectories in such a steady state, with the trap moving at a speed v_1 , and we then changed the speed from v_1 to v_2 over a time interval of duration τ , after which we continued to move the trap at speed v_2 (Fig. 2a). Thus, the trap speed v played the role of the parameter α in our earlier discussion. Modeling the bead as an overdamped Brownian particle (18), the steady-state distribution is $\rho_{ss}(x; v) \propto \exp[-\beta(\kappa x + \gamma v)^2/2\kappa]$, where x is the bead displacement relative to the minimum of the trap and β is the inverse temperature of the solution. Eq. 2 then gives us $Y = (\beta\gamma/\kappa) \int_0^\tau dt \dot{v}(t) [\kappa x(t) + \gamma v(t)]$, where $x(t)$ specifies the motion of the bead during a given realization of the process, and $v(t)$ is the (externally imposed) trap speed.

specifies a unit of power relevant to our dragged particle, just as β^{-1} specifies the relevant unit of energy in the equilibrium case. Last, in Eqs. 5, the work W is the contribution to the net change in the internal energy of the system, E , arising from increments in the value of α (25–27). Thus, by drawing parallels between $\bar{P}$ and the free energy ($\bar{P}/2 \sim F$), q and temperature ($q \sim \beta^{-1}$), and $\Delta_v P$ and mechanical work ($-\Delta_v P \sim W$), the analogy between equilibrium laws of Eqs. 5 and the nonequilibrium laws of Eqs. 7 is complete.

Our experiments support the suggestion that transitions between nonequilibrium steady states are subject to laws similar to those governing transitions between equilibrium states (Eqs. 5). These findings motivate several questions. First, does a steady-state thermodynamic formalism support a zeroth law? Such a law would define an effective temperature of nonequilibrium steady states (28). Second, can Eq. 4 also be understood in terms of the monotonic growth of some global quantity, just as the equilibrium second law states that the combined entropy of a system and its thermal surroundings never decreases? Last, at the microscopic level, can nonequilibrium steady states be described

by a universal theory for ρ_{ss} , analogous to the Boltzmann–Gibbs distribution for equilibrium states? If so, this result would greatly increase the predictive power of Eq. 4 as a “generalized second law” because the microscopic nature of the steady state enters into the definition of the quantity Y by means of the function $\phi = -\ln \rho_{ss}$. For now, we can apply Eqs. 3 and 4 only to systems for which ρ_{ss} can be deduced from a simple model or from direct experimental measurement of the distribution. Theoretical developments along these lines should be readily testable with the use of microscopic systems such as those described here, whose fluctuations are directly measurable.

We thank Yoshitsugu Oono, Ignacio Tinoco, Jr., Hari Shroff, Nathan Clack, and Merek Siu for valuable discussions and suggestions. J.L. is supported by the Director’s Fellowship of Lawrence Berkeley National Laboratory. This work was supported by National Institutes of Health Grant GM-32543; U.S. Department of Energy Grants DE-AC03-76SF00098 and W-7405-ENG-36; National Science Foundation Grants MBC-9118482 and DBI-9732140; a David and Lucile Packard Foundation grant (to C.J.B.); and the Spanish Ministerio de Ciencia y Tecnología Grant BFM2001-3525.

1. Chow, W. W., Scheider, H. C., Koch, S. W., Chang, C., Chrostowski, L. & Chang-Hasnain, C. J. (2002) *IEEE J. Quantum Electron.* **38**, 402–409.
2. Qian, H., Saffarian, S. & Elson, E. L. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 10376–10381.
3. Goldbeter, A. (1997) in *Biochemical Oscillations and Cellular Rhythms: The Molecular Bases of Periodic and Chaotic Behaviour* (Cambridge Univ. Press, London).
4. Wang, H. & Oster, G. (1998) *Nature* **396**, 279–282.
5. Keizer, J. (1987) in *Statistical Thermodynamics of Nonequilibrium Processes* (Springer, Berlin).
6. Jou, D., Casas-Vázquez, J., and Lebon, G. (1988) *Rep. Prog. Phys.* **51**, 1105–1179.
7. Eu, B. C. (1998) in *Nonequilibrium Statistical Mechanics* (Kluwer Academic, Dordrecht, The Netherlands).
8. Oono, Y. & Paniconi, M. (1998) *Prog. Theor. Phys. Suppl.* **130**, 29–44.
9. Finn, C. B. P. (1993) in *Thermal Physics* (Chapman and Hall, London), 2nd Ed..
10. Keizer, J. (1985) *J. Chem. Phys.* **82**, 2751–2771.
11. Hatano, T. & Sasa, S. (2001) *Phys. Rev. Lett.* **86**, 3463–3466.
12. Chandler, D. (1987) in *Introduction to Modern Statistical Mechanics* (Oxford Univ. Press, New York), Sect. 5.5.
13. Fauchaux, L. P., Bourdieu, L. S., Kaplan, P. D. & Libchaber, A. J. (1995) *Phys. Rev. Lett.* **74**, 1504–1507.
14. Hales, J., Zhukov, A., Roy, R. & Dykman, M. I. (2000) *Phys. Rev. Lett.* **85**, 78–81.
15. Wang, G. M., Seveick, E. M., Mittag, E., Searls, D. J. & Evans, D. J. (2002) *Phys. Rev. Lett.* **89**, 50601–50604.
16. Hummer, G. & Szabo, A. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 3658–3661.
17. Liphardt, J., Dumont, S., Smith, S. B., Tinoco, I., Jr., & Bustamante, C. (2002) *Science* **296**, 1832–1835.
18. Mazonka, O. & Jarzynski, C. (Dec. 7, 1999) arXiv.org:cond-mat 9912121.
19. Smith, S.B., Cui, Y. & Bustamante, C. (2003) *Methods Enzymol.* **361**, 134–162.
20. Gore, J., Ritort, F. & Bustamante, C. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 12564–12569.
21. Zuckerman, D. M. & Woolf, T. B. (2002) *Chem. Phys. Lett.* **351**, 445–453.
22. Zuckerman, D. M. & Woolf, T. B. (2002) *Phys. Rev. Lett.* **89**, 180602–180604.
23. Park, S., Khalili-Araghi, F., Tajkhorshid, E. & Schulten, K. (2003) *J. Chem. Phys.* **119**, 3559–3566.
24. Ritort, F., Bustamante, C., Tinoco, I. Jr. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 13544–13548.
25. Jarzynski, C. (1997) *Phys. Rev. Lett.* **78**, 2690–2693.
26. Jarzynski, C. (1997) *Phys. Rev. E* **56**, 5018–5035.
27. Crooks, G. E. (1998) *J. Stat. Phys.* **90**, 1481–1487.
28. Casas-Vázquez, J. & Jou, D. (2003) *Rep. Prog. Phys.* **66**, 1937–2023.
29. Efron, B. & Tibshirani, R. J. (1993) in *An Introduction to the Bootstrap* (CRC, New York).